

(FILE 'HOME' ENTERED AT 18:09:08 ON 06 SEP 2002)

FILE 'BIOSIS, MEDLINE, INPADOC, CAPLUS' ENTERED AT 18:09:22 ON 06 SEP 2002

L1 3 BONE CELL SPHEROID  
L2 49 BONE AND SPHEROID  
L3 46 L2 NOT L1  
L4 32 DUPLICATE REMOVE L3 (14 DUPLICATES REMOVED)  
L5 83937 CULTUR? AND (BONE OR OSTEO?)  
L6 3521 L5 AND (TRANSFORMING GROWTH FACTOR#)  
L7 238 L6 AND (SERUM FREE)  
L8 6 L7 AND (BONE PRECURSOR CELL#)  
L9 55 L7 AND OSTEOBLAST#  
L10 30 DUPLICATE REMOVE L9 (25 DUPLICATES REMOVED)  
L11 3135 (SERUM FREE MEDI##) AND ITS  
L12 3135 (SERUM FREE MEDI##) AND "ITS+"  
L13 124 (SERUM FREE MEDI##) (5A) "ITS+"  
L14 60 DUPLICATE REMOVE L13 (64 DUPLICATES REMOVED)

=>

AN 1995:349833 CAPLUS

DN 122:123874

TI Regulation of human **bone** marrow-derived **osteoprogenitor** cells by **osteogenic** growth factors

AU Long, Michael W.; Robinson, J. A.; Ashcraft, E. A.; Mann, Kenneth G.

CS Dep. Pediatr., Univ. Michigan, Ann Arbor, MI, 48109, USA

SO Journal of Clinical Investigation (1995), 95(2), 881-7

CODEN: JCINAO; ISSN: 0021-9738

PB Rockefeller University Press

DT Journal

LA English

AB Human **bone** marrow contains a distinct cell population that expresses **bone** proteins and responds to **transforming growth factor .beta.1** (TGF-.beta.), but not to hematopoietic growth factors (Long, M. W. et al, 1990). The authors now report the isolation, characterization, and growth factor responsiveness of these precursors to human **osteoblasts** and the identification of a human **osteoprogenitor** cell. Immunol. sepn. of human **bone** marrow nonadherent low-d. (NALD) cells results in a marked enrichment of cells that express **osteocalcin**, **osteonectin**, and **bone** alk. phosphatase. Flow cytometric anal. show that distinct cell subpopulations exist among these isolated cells. The majority of the **bone** antigen-pos. cells are approx. the size of a lymphocyte, whereas other, less frequent antibody-sepd. subpopulations consist of **osteoblast**-like cells and **osteoprogenitor** cells. In **serum-free** cultures, TGF-.beta. stimulates the small, antigen-pos. cells to become **osteoblast**-like, as these cells both increase in size, and express increased levels of **osteocalcin** and alk. phosphatase. Antibody-sepd. cells also contain a sep. population of clonal progenitor cells that form colonies of **osteoblast**-like cells when cultured in **serum-free**, semi-solid media. Two types of human **osteoprogenitor** cells are obsd.: a colony-forming cell (CFC) that generates several hundred **bone** antigen-pos. cells, and more mature cluster-forming cell that has a lesser proliferative potential and thus generates clusters of 20-50 antigen-pos. cells. **Osteopoietic** colony-forming cells and cluster-forming cells have an obligate but differential requirement for **osteogenic** growth factors. The CFCs respond to TGF-.beta., basic fibroblast growth factor (bFGF), **bone** morphogenic protein-2 (BMP-2), and 1,25-dihydroxyvitamin D3 (1,25-OH D3). In contrast to the colony-forming cells, cluster-forming cells are regulated predominantly by 1,25-OH D3 and TGF-.beta., but fail to respond to bFGF. The authors conclude that human **bone** marrow contains a nonhematogenous, heterogeneous population of **bone precursor** cells among which exists a population of proliferating **osteoprogenitor** cells. Further characterization of these **bone precursor** cell populations should yield important information on their role in **osteogenesis** in both health and disease.

=&gt;

L1 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2002 ACS  
AN 2000:628798 CAPLUS  
DN 133:279465  
TI Three-dimensional cellular development is essential for ex vivo formation  
of human bone  
AU Kale, Sujata; Biermann, Sybil; Edwards, Claire; Tarnowski, Catherine;  
Morris, Michael; Long, Michael William  
CS Department of Pediatrics, Surgery, Chemistry, and Comprehensive Cancer  
Center, University of Michigan, Ann Arbor, MI, 48109, USA  
SO Nature Biotechnology (2000), 18(9), 954-958  
CODEN: NABIF9; ISSN: 1087-0156  
PB Nature America Inc.  
DT Journal  
LA English  
AB Tissue engineering of human bone is a complex process, as the functional  
development of bone cells requires that regulatory signals be temporally  
and spatially ordered. The role of three-dimensional cellular  
interactions is well understood in embryonic osteogenesis, but in vitro  
correlates are lacking. Here we report that in vitro serum-free  
transforming growth factor (TGF)- $\beta$ 1 stimulation of osteogenic cells  
immediately after passage results in the formation of three-dimensional  
cellular condensations (bone cell spheroids) within 24 to 48 h. In turn,  
**bone cell spheroid** formation results in the  
up-regulation of several bone-related proteins (e.g., alk. phosphatase,  
type 1 collagen, osteonectin) during days 3-7, and the concomitant  
formation of micro-cryst. bone. This system of ex vivo bone formation  
should provide important information on the physiol., biol. and mol. basis  
of osteogenesis.  
RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=>

L3 ANSWER 179 OF 181 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
DUPLICATE 116  
AN 1978:194198 BIOSIS  
DN BA66:6695  
TI **BONE CELLS A SERUM-FREE MEDIUM SUPPORTS**  
**PROLIFERATION IN PRIMARY CULTURE.**  
AU BURKS J K; PECK W A  
CS DEP. MED., JEW. HOSP. ST. LOUIS, WASH. UNIV. SCH. MED., ST. LOUIS, MO.  
63110, USA.  
SO SCIENCE (WASH D C), (1978) 199 (4328), 542-544.  
CODEN: SCIEAS. ISSN: 0036-8075.  
FS BA; OLD  
LA English  
AB Bone cells isolated from embryonic rat calvaria increase in number  
2-3-fold when cultured at high, but not at low, population densities in a  
serum-free medium that contains albumin. Cultured cells respond to  
parathyroid hormone and exhibit a marked rise in alkaline phosphatase  
activity during proliferation, which suggests the progressive  
differentiation or preferential growth of osteoblast-like cells.

# Bone Cells: A Serum-Free Medium Supports Proliferation in Primary Culture

**Abstract.** Bone cells isolated from embryonic rat calvaria increase in number two- to threefold when cultured at high, but not at low, population densities in a serum-free medium that contains albumin. Cultured cells respond to parathyroid hormone and exhibit a marked rise in alkaline phosphatase activity during proliferation, which suggests progressive differentiation or preferential growth of osteoblast-like cells.

Studies in which isolated bone cells in primary culture are used have advanced our understanding of the actions of major bone-seeking hormones (1). Cultured cells are sensitive to near physiological concentrations of several hormones and retain the capacity to perform differentiated functions (2, 3). A strict requirement for serum to support the growth of bone cells in vitro, however, has impeded further clarification of the regulation of cell proliferation and differentiation, since serum contains defined and undefined growth-modifying substances (4). Attempts to grow non-skeletal cells in the absence of serum have met with but limited success. Additions of specific nutrients, hormones, or polypeptides reduce, but do not eliminate, serum requirements for growth of certain cell types in primary culture (5). Although several established cell lines have been found to grow in serum-free medium, they are less suitable than primary cultures for studies of differentiated functions (5). We report the development of a serum-free medium that supports the proliferation of isolated bone cells in primary culture.

Bone cells were isolated from the frontal and parietal bones of rat fetuses 19 to 21 days old, as previously described (1). Cartilage and periosteal connective tissue were removed by sharp dissection prior to enzymatic digestion with crude collagenase (1). Cultures were seeded at an initial density of  $10^5$  cells ( $10^5$  cell/cm<sup>2</sup>) per Falcon petri dish (diameter, 35 mm) and were maintained in a humid atmosphere of 2 percent CO<sub>2</sub> and 98 percent air. Cells were grown in the Fitton-Jackson BGJ<sub>1</sub> medium (6), previously used to study bone cell metabolism (7), and modified so that its ionic composition resembled that of the bone fluid as described by Neuman and Ramp (8) (Table 1). The sodium and potassium concentrations were 125 mM and 25 mM, respectively. This medium (2.0 ml) was replaced three times a week. Cell number (after trypsinization), alkaline phosphatase activity, and DNA content were estimated on days 1, 5, 9, and 12 of culture (Fig. 1). Replicate cell cultures were exposed to [<sup>3</sup>H]thymidine (1.0  $\mu$ Ci per 2.0 ml of culture medium) (9) for 1 hour, and the radioactivity in the acid-insoluble fraction (10) was estimated

with a Packard Tri-Carb liquid-scintillation spectrophotometer.

Cells began to attach to the plastic surface within 5 minutes of seeding. Although 35 percent of the inoculated cells remained firmly attached after the first 24 hours, failure to attach was not related to cell death, since 75 percent of the unattached cells were viable as judged by trypan blue exclusion. In view of the well-known limitations of dye exclusion methods in assessing cell integrity, the unattached cells may have been damaged sufficiently to prevent attachment. Alternatively, the attached cells may represent a different type of cell than the remainder of the aggregate cell harvest. Microscopic inspection of the culture plates suggested cell proliferation by day 5. This impression was confirmed by the presence of statistically significant increases in the cell number and total DNA that continued through day 9 (Fig. 1). The incorporation of [<sup>3</sup>H]thymidine rose progressively to a maximum of  $42,160 \pm 2,232$  disintegrations per minute (dpm) per culture (mean  $\pm$  standard error of the mean) on day 7; then gradually fell to  $17,237 \pm 1,281$  dpm per culture on day 12, by which time the cultures were nearly confluent. The conditions used in these experiments consistently supported bone cell proliferation. Figure 1 presents data accumulated from six separate experiments. Bone cells cultured at lower initial population densities ( $5 \times 10^4$  cells per square centimeter) failed to proliferate. Instead there was gradual cell attrition over a 12-day period.

Three lines of evidence suggest the





Fig. 1 (left). Growth of isolated subperiosteal bone cells seeded at an initial population density of  $10^5$  cell/cm<sup>2</sup>. Each point represents the mean and standard error of data pooled from six separate experiments. Day 0 represents the initial inoculation; all other points represent firmly attached cells removed from the plates after trypsinization. Solid line is cell number  $\times 10^5$  per culture; dashed line, alkaline phosphatase activity [(micromoles of *p*-nitrophenol per hour)  $\times 10^3$ ]; dotted line, DNA in micrograms per culture. Arrows represent the days on which the medium was replaced. Following enzymatic digestion of the calvaria, the dispersed cells were rinsed four times with isolation medium, allowed to stand at room temperature for 5 minutes in isotonic ammonium chloride to lyse red blood cells (22), and then washed and suspended in isolation medium for hemocytometer counting. Alkaline phosphatase activity was measured by the method of Koyama and Ono (23) at a pH of 10.3. DNA was determined by the indole method of Hubbard *et al.* (24), in which hydrolysis for 20 minutes at 90°C and a volume of 1.0 ml was used, or by the diphenylamine method of Leyra and Kelley (25). Both methods gave identical results when corrected for recovery of internal standards. Fig. 2 (right). Morphological characteristics of bone cells cultured for 12 days in bone fluid medium. (a) Phase-contrast photomicrograph demonstrating dense aggregates of cells in a near confluent monolayer. (b) Cytoplasmic and nuclear alkaline phosphatase activity. Cultures were fixed at 4°C (30 seconds) with 10 percent formalin in methanol and stained by the method of Ackerman (11). Inset represents the region where cells form dense aggregates. Cells devoid of alkaline phosphatase activity are not visible.

medium was replaced. Following enzymatic digestion of the calvaria, the dispersed cells were rinsed four times with isolation medium, allowed to stand at room temperature for 5 minutes in isotonic ammonium chloride to lyse red blood cells (22), and then washed and suspended in isolation medium for hemocytometer counting. Alkaline phosphatase activity was measured by the method of Koyama and Ono (23) at a pH of 10.3. DNA was determined by the indole method of Hubbard *et al.* (24), in which hydrolysis for 20 minutes at 90°C and a volume of 1.0 ml was used, or by the diphenylamine method of Leyra and Kelley (25). Both methods gave identical results when corrected for recovery of internal standards. Fig. 2 (right). Morphological characteristics of bone cells cultured for 12 days in bone fluid medium. (a) Phase-contrast photomicrograph demonstrating dense aggregates of cells in a near confluent monolayer. (b) Cytoplasmic and nuclear alkaline phosphatase activity. Cultures were fixed at 4°C (30 seconds) with 10 percent formalin in methanol and stained by the method of Ackerman (11). Inset represents the region where cells form dense aggregates. Cells devoid of alkaline phosphatase activity are not visible.

preferential growth or differentiation (or both) of osteoblast-like cells in this culture system. (i) A marked rise in alkaline phosphatase activity per cell accompanied cell growth (Fig. 1). (ii) The number of cells that contained prominent cytoplasmic alkaline phosphatase activity increased from less than 1 percent on day 1 to more than 25 percent on day 12 of culture (Fig. 2). (iii) Cells cultured for 9 to 12 days responded to parathyroid hormone [synthetic bovine parathyroid hormone, amino acids 1-34 (11, 12), 100 ng/ml for 5.0 minutes] with a tenfold increase in 3',5'-adenosine monophosphate (cyclic AMP) when treated as described (2) (data not shown). There is every reason to believe that osteoblasts contain copious amounts of cytoplasmic alkaline phosphatase (13). Luben *et al.* (14) have provisionally characterized as osteoblasts those bone cells that are rich in alkaline phosphatase and respond to parathyroid hormone with large increases in cyclic AMP. The observed increase in alkaline phosphatase activity might reflect recovery from damage to surface enzyme caused by exposure of the bone cell to collagenase during isolation, rather than de novo differentiation. In favor of the latter interpretation, however, is the finding of Miedema (15) that the loss of alkaline phosphatase activity which accompanied collagenase treatment of an established cell line was not reversible. Definitive identification of the surviving cells as osteoblasts will require the demonstration that they are capable of producing bone.

Miedema (15) has demonstrated a similar density-dependent increase in alkaline phosphatase activity in cultured human HEp2 and WISH cells, as have Binderman *et al.* (7) in an isolated bone cell system. In each case, the incubation medium was supplemented with fetal calf serum. Yamane *et al.* (16) have initiated primary cultures of ascites tumor cells in a defined medium supplemented with 1 percent fat-containing serum albumin. In addition to providing essential fats, albumin may have served to trap toxic substances (17). Fat-poor albumin, such as that used in our system, was not found to support the growth of the ascites cells (16).

To determine whether the unusual ionic composition of the incubation medium used in our experiments (125 mM sodium, 25 mM potassium) influenced cell growth and apparent differentiation, we examined the effect of a medium containing more physiological concentrations of sodium (145 mM) and potassium (5 mM). Direct comparison of these media revealed no differences in plating efficiency

or in the rate of cell proliferation. Cells cultured in the standard medium, however, exhibited only 50 percent of the maximum alkaline phosphatase activity and were only 50 percent as responsive to parathyroid hormone (see above) as those grown in the low sodium-high potassium medium (data not shown). No proliferation was observed in medium containing 155 mM sodium

and 25 mM potassium (mean osmolality, 385 milliosmoles). Hence proliferation may be sensitive to osmolality, and apparent differentiation to the concentrations of sodium and potassium.

Isolated periosteal cells, removed from the superior and inferior periosteal surfaces by sharp dissection prior to collagenase digestion (18), failed to survive in the serum-free medium even when

Table 1. Comparison of Fitton-Jackson BGJ<sub>h</sub> with bone fluid medium. Fitton-Jackson modified BGJ<sub>h</sub> medium (Gibco) has a mean osmolality of 390 milliosmoles. To prepare the bone fluid medium, the salts and organic components are dissolved in 6 to 7 dl of glass-distilled deionized water. One deciliter of commercially prepared BGJ<sub>h</sub> (Fitton-Jackson modified) amino acids and vitamins (10 ×) without ascorbic acid and glutamine is added (Gibco, quotation No. 8202) followed by 1000 mg of Pentex albumin (Miles Laboratories). The pH is adjusted to 7.45 at room temperature with 0.4M NaOH and the volume adjusted to 1 liter; NaHCO<sub>3</sub> is added prior to membrane sterilization. The final pH is 7.55, which is optimum for cell attachment and collagen biosynthesis (21). The mean osmolality of the bone fluid medium is 350 milliosmoles. The isolation medium is prepared in a similar manner, but has a final pH of 7.00.

Component	Concentration	
	Fitton-Jackson BGJ <sub>h</sub> (mg/liter) (mole/liter)	Bone fluid medium (mole/liter) (mg/liter)
CaCl <sub>2</sub> · 2H <sub>2</sub> O		73.52 (5.0 × 10 <sup>-4</sup> )
Calcium lactate	555.00 (1.8 × 10 <sup>-3</sup> )	
MgSO <sub>4</sub> · 7H <sub>2</sub> O	200.00 (8.1 × 10 <sup>-4</sup> )	98.60 (4.0 × 10 <sup>-4</sup> )
K <sub>2</sub> HPO <sub>4</sub>		261.28 (1.5 × 10 <sup>-3</sup> )
KH <sub>2</sub> PO <sub>4</sub>	160.00 (1.2 × 10 <sup>-3</sup> )	
NaH <sub>2</sub> PO <sub>4</sub> · H <sub>2</sub> O	90.00 (6.5 × 10 <sup>-4</sup> )	41.40 (3.0 × 10 <sup>-4</sup> )
NaCl	5,300.00 (9.1 × 10 <sup>-2</sup> )	5,844.00 (1.0 × 10 <sup>-1</sup> )
KCl	400.00 (5.4 × 10 <sup>-3</sup> )	1,640.26 (2.2 × 10 <sup>-2</sup> )
NaHCO <sub>3</sub>	3,500.00 (4.2 × 10 <sup>-2</sup> )	1,260.60 (1.5 × 10 <sup>-2</sup> )
FeSO <sub>4</sub> · 7H <sub>2</sub> O		0.83 (3.0 × 10 <sup>-5</sup> )
Hepes†		4,766.00 (2.0 × 10 <sup>-2</sup> )
D-Glucose	10,000.00 (5.5 × 10 <sup>-2</sup> )	2,000.00 (1.1 × 10 <sup>-2</sup> )
Sodium acetate (3H <sub>2</sub> O)	50.00 (6.0 × 10 <sup>-4</sup> )	54.43 (4.0 × 10 <sup>-4</sup> )
α-Ketoglutaric acid		29.22 (2.0 × 10 <sup>-4</sup> )
Phenol red	20.00 (5.6 × 10 <sup>-5</sup> )	5.00 (1.4 × 10 <sup>-5</sup> )
L-Alanine	250.00 (2.8 × 10 <sup>-3</sup> )	250.00 (2.8 × 10 <sup>-3</sup> )
L-Arginine	175.00 (1.0 × 10 <sup>-3</sup> )	175.00 (1.0 × 10 <sup>-3</sup> )
L-Aspartic acid	150.00 (1.1 × 10 <sup>-3</sup> )	150.00 (1.1 × 10 <sup>-3</sup> )
L-Cysteine HCl · H <sub>2</sub> O	90.00 (5.1 × 10 <sup>-4</sup> )	90.00 (5.1 × 10 <sup>-4</sup> )
L-Glutamine	200.00 (1.4 × 10 <sup>-3</sup> )	292.3 (2.0 × 10 <sup>-3</sup> )
Glycine	800.00 (1.1 × 10 <sup>-2</sup> )	800.00 (1.1 × 10 <sup>-2</sup> )
L-Histidine	150.00 (9.6 × 10 <sup>-4</sup> )	150.00 (9.6 × 10 <sup>-4</sup> )
L-Isoleucine	30.00 (2.3 × 10 <sup>-4</sup> )	30.00 (2.3 × 10 <sup>-4</sup> )
L-Leucine	50.00 (3.8 × 10 <sup>-4</sup> )	50.00 (3.8 × 10 <sup>-4</sup> )
L-Lysine	240.00 (1.3 × 10 <sup>-3</sup> )	240.00 (1.3 × 10 <sup>-3</sup> )
L-Methionine	50.00 (3.4 × 10 <sup>-4</sup> )	50.00 (3.4 × 10 <sup>-4</sup> )
L-Phenylalanine	150.00 (3.0 × 10 <sup>-4</sup> )	150.00 (3.0 × 10 <sup>-4</sup> )
L-Proline	400.00 (3.5 × 10 <sup>-3</sup> )	400.00 (3.5 × 10 <sup>-3</sup> )
L-Serine	200.00 (1.9 × 10 <sup>-3</sup> )	200.00 (1.9 × 10 <sup>-3</sup> )
L-Threonine	75.00 (6.3 × 10 <sup>-4</sup> )	75.00 (6.3 × 10 <sup>-4</sup> )
L-Tryptophan	40.00 (2.0 × 10 <sup>-4</sup> )	40.00 (2.0 × 10 <sup>-4</sup> )
L-Tyrosine	40.00 (2.2 × 10 <sup>-4</sup> )	40.00 (2.2 × 10 <sup>-4</sup> )
Dl-Valine	65.00 (5.6 × 10 <sup>-4</sup> )	65.00 (5.6 × 10 <sup>-4</sup> )
α-Tocopherol phosphate	1.00 (2.3 × 10 <sup>-6</sup> )	1.00 (2.3 × 10 <sup>-6</sup> )
Ascorbic acid	50.00 (2.8 × 10 <sup>-4</sup> )	50.00 (2.8 × 10 <sup>-4</sup> )
Biotin	0.20 (8.2 × 10 <sup>-7</sup> )	0.20 (8.2 × 10 <sup>-7</sup> )
Calcium pantothenate	0.20 (4.2 × 10 <sup>-7</sup> )	0.20 (4.2 × 10 <sup>-7</sup> )
Choline chloride	50.00 (3.6 × 10 <sup>-4</sup> )	50.00 (3.6 × 10 <sup>-4</sup> )
Folic acid	0.20 (4.5 × 10 <sup>-7</sup> )	0.20 (4.5 × 10 <sup>-7</sup> )
i-Inositol	0.20 (1.1 × 10 <sup>-6</sup> )	0.20 (1.1 × 10 <sup>-6</sup> )
Nicotinamide	20.00 (1.6 × 10 <sup>-4</sup> )	20.00 (1.6 × 10 <sup>-4</sup> )
p-Amino benzoic acid	2.00 (1.6 × 10 <sup>-5</sup> )	2.00 (1.6 × 10 <sup>-5</sup> )
Pyridoxal phosphate	0.20 (8.1 × 10 <sup>-7</sup> )	0.20 (8.1 × 10 <sup>-7</sup> )
Riboflavin	0.20 (5.3 × 10 <sup>-7</sup> )	0.20 (5.3 × 10 <sup>-7</sup> )
Thiamine-HCl	4.00 (1.2 × 10 <sup>-5</sup> )	4.00 (1.2 × 10 <sup>-5</sup> )
Vitamin B <sub>12</sub>	0.04 (2.9 × 10 <sup>-6</sup> )	0.04 (2.9 × 10 <sup>-6</sup> )

\*The amount of NaCl added depends upon the NaCl content of the commercially obtained amino acid-vitamin mixture. The final ion concentrations are similar to the bone fluid described by Neuman and Ramp: Na<sup>+</sup>, 125 mM; K<sup>+</sup>, 25 mM; Ca<sup>2+</sup>, 0.5 mM; Mg<sup>2+</sup>, 0.4 mM; Cl<sup>-</sup>, 130 mM; and P<sub>i</sub>, 1.8 mM (8). †N-2-Hydroxyethylpiperazine-N'-2-ethanesulfonic acid. ‡Anhydrous sodium acetate.

seeded at high initial population densities ( $10^5$  cell/cm<sup>2</sup>). Seventeen percent of the cells attached, but only 5 percent remained by day 5. The alkaline phosphatase activity was low ( $28.3 \pm 5.3$   $\mu$ mole per h ur per culture) in comparison with that of cell cultures prepared from subperiosteal bone. Fibroblasts obtained by collagenase digestion of fetal rat skin died over the initial 2- to 3-day period in culture.

Growth and apparent differentiation at a high, but not at a low, initial population density may reflect the continuing action of tissue or serum growth factors, or both, that adhered to the cells during preparation. Alternate explanations include (i) the elaboration by cultured cells of their own growth factors or essential nutrients that are absent from the medium (19) and (ii) inactivation by the larger cell mass of toxic substances in the medium (20). The latter possibility would explain not only the failure of cells to proliferate when cultured at low density but also the low plating efficiency. Although there was appreciable cell death during the first 24 hours in culture, stimulation of proliferation cannot be attributed easily to growth-promoting factors released into the incubation medium by the dying cells. The incubation medium was replaced completely at 24 hours (Fig. 1), whereas proliferation was evident long afterward. Moreover, fibroblasts and periosteal cells did not proliferate, despite greater initial cell attrition. The system described herein should permit clarification of the mechanism of density-dependent proliferation and an examination of the effects of growth factors on the proliferation and differentiation of bone cells in vitro in the absence of complex and ill-defined serum additives.

JAMES K. BURKS  
WILLIAM A. PECK

Department of Medicine,  
Jewish Hospital of St. Louis,  
Washington University School of  
Medicine, St. Louis, Missouri 63110

#### References and Notes

- W. A. Peck, S. J. Birge, Jr., S. A. Fedak, *Science* 146, 1476 (1964).
- W. A. Peck, J. Carpenter, K. Messinger, D. DeBra, *Endocrinology* 92, 692 (1973).
- W. A. Peck, K. Messinger, J. Brandt, J. Carpenter, *J. Biol. Chem.* 244, 4174 (1969); S. B. Rodan and G. A. Rodan, *ibid.* 249, 3068 (1974); G. L. Wong and D. V. Cohn, *Proc. Natl. Acad. Sci. U.S.A.* 72, 3167 (1975); G. Wong and D. Cohn, *Nature (London) New Biol.* 252, 713 (1975).
- H. M. Temin, R. W. Pierson, N. C. Dulak, in *Growth, Nutrition and Metabolism of Cells in Culture*, G. H. Rothblat and J. Cristofalo, Eds. (Academic Press, New York, 1972), p. 49; R. W. Holley and J. A. Kiernan, *Proc. Natl. Acad. Sci. U.S.A.* 63, 300 (1968).
- R. W. Pumper, *Science* 128, 363 (1958); D. V. Young, *J. Cell. Physiol.* 89, 133 (1976); V. J. Evans, J. C. Bryant, H. A. Kerr, E. L. Schilling, *Exp. Cell Res.* 36, 439 (1964); L. T. Dupree, K. K. Sanford, B. B. Westfall, A. B. Covalsky, *ibid.* 28, 381 (1962); H. Eagle, V. J. Oyama, K. A. Piez, *J. Biol. Chem.* 235, 1719 (1960); J. R. Birch and S. J. Pirt, *J. Cell Sci.* 5, 132 (1969); W. L. McKeethan, W. G. Hamilton, R. G. Ham, *Proc. Natl. Acad. Sci. U.S.A.* 73, 2023 (1976); K. K. Sanford, L. T. Dupree, A. B. Covalsky, *Exp. Cell Res.* 31, 345 (1963); D. Gospodarowicz and J. S. Moran, *Annu. Rev. Biochem.* 45, 531 (1976); R. W. Holley, *Nature (London)* 258, 487 (1975); K. Higuchi and R. C. Robinson, *In Vitro* 9 (No. 2), 114 (1973); I. Hayashi and G. H. Sato, *Nature (London)* 259, 132 (1976).
- J. D. Biggers, R. B. L. Gwatkin, S. Heyner, *Exp. Cell Res.* 26, 41 (1961); for formulation of the Fitton-Jackson modification, see catalog of Grand Island Biological Co.
- L. G. Raisz and I. Niemann, *Endocrinology* 85, 446 (1967); I. Binderman, D. Duksin, A. Harell, E. Kalz, L. Sachs, Jr., *J. Cell Biol.* 61, 427 (1974).
- W. F. Neuman and W. K. Ramp, in *Cellular Mechanisms for Calcium Transfer and Homeostasis*, G. Nichols, Jr., and R. H. Wasserman, Eds. (Academic Press, New York, 1971), p. 197.
- [methyl-<sup>3</sup>H]Thymidine, 20 c/mole (New England Nuclear).
- S. P. Nissley, J. Passamani, P. Short, *J. Cell. Physiol.* 89, 393 (1976).
- G. A. Ackermann, *Lab. Invest.* 11, 563 (1962).
- Parathyroid hormone, residues 1-34 (Beckman).
- J. J. Pritchard, in *The Biochemistry and Physiology of Bone*, G. H. Bourne, Ed. (Academic Press, New York, 1956), p. 179; M. S. Burstone, in *Calcification in Biological Systems*, R. F. Sognnaes, Ed. (AAAS, Washington, D.C., 1960), p. 217.
- R. A. Luben, G. L. Wong, D. V. Cohn, *Endocrinology* 99, 526 (1976).
- E. Miedema, *Exp. Cell Res.* 53, 488 (1968).
- I. Yamane, D. Murakami, M. Kato, *Proc. Soc. Exp. Biol. Med.* 149, 439 (1975).
- J. Michl, *Exp. Cell Res.* 23, 324 (1961); M. Moskowitz and D. M. Schenck, *ibid.* 38, 523 (1965).
- W. A. Peck, J. K. Burks, J. Wilkins, S. B. Rodan, G. A. Rodan, *Endocrinology* 100, 1357 (1977).
- H. Eagle and K. A. Piez, *J. Exp. Med.* 116, 29 (1962).
- R. G. Ham, *In Vitro* 10, 119 (1974).
- T. P. Nigra, G. R. Martin, N. Cagle, *Biochem. Biophys. Res. Commun.* 53, 272 (1973); R. S. Gardner, *J. Cell Biol.* 42, 320 (1969); H. Eagle, *Science* 174, 500 (1971).
- W. Boyle, *Transplantation* 6, 761 (1968).
- H. Koyama and T. Ono, *Biochim. Biophys. Acta* 264, 497 (1972).
- R. W. Hubbard, W. T. Matthew, D. A. Dubowik, *Anal. Biochem.* 38, 190 (1970); R. W. Hubbard, W. T. Matthew, D. W. Moulton, *ibid.* 46, 461 (1972).
- A. Leyra, Jr., and W. N. Kelley, *ibid.* 62, 1973 (1974).
- We thank J. Weil and I. Dowling for their technical assistance. This work was supported by research grant AM-19853 from the National Institute of Arthritis, Metabolism, and Digestive Diseases.

31 May 1977; revised 19 August 1977

## Cyclopropanecarboxylic Acid: Chain Elongation to $\omega$ -Cyclopropyl Fatty Acids by Mammals and Plants

**Abstract.** Rats dosed orally with [carboxyl-<sup>14</sup>C]cyclopropanecarboxylic acid (or its hexadecyl ester) retain radioactivity in tissue as novel triacylglycerols. The most abundant <sup>14</sup>C-labeled metabolites were identified by gas-liquid chromatography-mass spectrometry as 13-cyclopropyltridecanoic and 15-cyclopropylpentadecanoic acids. Similar  $\omega$ -cyclopropyl fatty acids are produced by beagle dogs and a lactating cow, as well as by apple and orange trees.

Certain esters of cyclopropanecarboxylic acid (CPCA) are selectively toxic to phytophagous Acarina (spider mites) without significant insecticidal activity (1, 2). We have studied the metabolic fate of the hexadecyl ester of CPCA (cycloprate), which is currently under commercial development as a miticide. We have identified unusual metabolites arising from the apparent entry of CPCA into pathways of fatty acid anabolism, with net additions of acetate to give a homologous series of  $\omega$ -cyclopropyl fatty acids.

Duncombe and Aising (3) studied the

metabolism of <sup>14</sup>C-labeled CPCA in rat tissue in vitro and postulated the formation of unsaturated  $\omega$ -cyclopropyl fatty acids, but were unable to identify these products. Linscott *et al.* (4, 5) have shown that plants can elongate the carbon chains of 2,4-dichlorophenoxyalkanoic (for example, acetic (2,4-D) and butyric) acids by insertion of pairs of methylene groups. The major metabolites of these acids were identified as resulting from addition of one to three acetate units (that is, two to six CH<sub>2</sub> groups), whereas we find addition of up to eight acetates to CPCA. We know of

Table 1. Quantitative abundance of  $\omega$ -cyclopropyl fatty acids, as a percent of the administered dose of [<sup>14</sup>C]cycloprate, in portions of various organisms that were analyzed (6).

$\omega$ -Cyclopropyl fatty acids*	Rat carcass	Cow milk	Dog carcass	Apple fruit	Orange fruit
8(5cPr): 0		0.2			
10(7cPr): 0		0.3			
12(9cPr): 0		0.3			
14(11cPr): 0	1.0	0.7	0.4		
16(13cPr): 0	9.9	1.6	6.5		
18(15cPr): 0	2.5	0.2	2.7	4.5	7.6
18(15cPr): 1				1.9	3.7
20(17cPr): 1					0.3

\*Structural abbreviations are explained in the legend to Fig. 1.

ANSWER 121 OF 181 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

DUPLICATE 74

AN 1990:153290 BIOSIS

DN BA89:80708

TI LONG-TERM ORGAN CULTURE OF EMBRYONIC CHICK FEMORA A SYSTEM FOR INVESTIGATING BONE AND CARTILAGE FORMATION AT AN INTERMEDIATE LEVEL OF ORGANIZATION.

AU ROACH H I

CS ACAD. ORTHOPAED. UNIT, SOUTHAMPTON UNIV., GEN. HOSP., CF 86, TREMONA RD., SOUTHAMPTON S09 4XY, UK.

SO J BONE MINER RES, (1990) 5 (1), 85-100.

CODEN: JBMREJ. ISSN: 0884-0431.

FS BA; OLD

LA English

AB Bone organ culture is an experimental system in which skeletal cells remain within their extracellular matrix but are removed from systemic influences. Femurs from 14-day-old chick embryos, which contain bone and cartilage matrix in approximately equal proportions, were cultured for up to 9 days in a **serum-free** medium. Cell proliferation, differentiation into chondrocytes and osteoblasts, formation of **bone** and cartilage matrix, and in vitro mineralization as well as bone and cartilage resorption were assessed using histologic and analytic methods. Particular attention was paid to the differences between cartilage and bone growth and to interpreting ~~analytic data in the light of histologic observations. The first 2 days of~~ culture represented an "adaptation" period, characterized by the release of intracellular enzymes into the culture medium, probably as a consequence of cell breakdown. Days 3-9 in culture represented a period of "steady growth" during which skeletal cells continued to multiply in the absence of fetal serum and to secrete large amounts of bone and cartilage matrix. De novo mineralization could be induced by Ca-.beta.-glycerophosphate, but calcium deposits in tissues other than bone and cartilage were also induced. Resorption of bone or cartilage matrix was virtually absent. Bone organ culture facilitates the study of bone and cartilage formation at an intermediate level of organization and thereby provides the necessary link between in vivo studies and investigations at the cellular level.